

Biomarkers of Premalignant Neoplasia and their use as Surrogate Endpoints in Clinical Trials of Chemopreventive Agents

Charles W. Boone

*Division of Cancer Prevention and Control, Chemoprevention Branch,
National Cancer Institute, National Institutes of Health, Bethesda, MD 20892*

WHAT IS CANER CHEMOPREVENTION?

Cancer chemoprevention is the prevention of clinical cancer by giving drugs or dietary constituents prior to or during the early phases of precancerous neoplasia, i.e., while the neoplastic process is still confined to the intraepithelial compartment and has not yet become invasive. Chemoprevention includes the prevention and treatment of genomic instability, the earliest alteration in the carcinogenic process, which begins in normal appearing epithelium prior to the occurrence of dysplasia and long before invasiveness. Genomic instability is a condition of the cellular DNA characterized by ever-expanding structural abnormalities and mutations; it is produced by exposure to carcinogens from the environment or by inherited abnormalities of DNA repair, such as those found in Fanconi's Anemia, Ataxia Telangiectasia, or Xeroderma Pigmentosum. The major criterion for the diagnosis of cancer is invasiveness, or impending invasiveness evidenced by full-thickness severe dysplasia (also called carcinoma in situ). Thus, the treatment of intraepithelial neoplasia with drugs can be viewed either as the chemoprevention of neoplastic progression from the intraepithelial, precancerous state to the invasive, cancerous state, or as the chemotherapy of intraepithelial neoplasia.

HOW DO CANCER CHEMOPREVENTIVE DRUGS WORK? ANS: BY SLOWING OR STOPPING NEOPLASTIC CLONAL EVOLUTION

What is Clonal Evolution?

Clonal evolution is the continuous appearance within a neoplastic cell population of mutant cells able to escape ambient growth control mechanisms and form clonal expansions which compete with each other on the basis of fastest growth rate^{1,2}. Vogelstein³ has revealed an outstanding demonstration of clonal evolution in adenomatous polyps of the colon: he showed that during the progression of intraepithelial neoplasia in the polyps, there occurs a series of genetic lesions, each of which is associated with a wave of overgrowing clonal cells.

Clonal evolution is the underlying mechanism by which neoplasms tend to develop ever-greater structural and functional diversity. The emergence of resistant clonal cell variants out of this diversity is the means by which neoplasms escape normal growth controls and also become resistant to chemotherapeutic agents and radiotherapy. Frequently, more than one genetically altered clone is found in different parts of a tumor, either by direct visualization or by genetic analysis. The different histological patterns used by the

pathologist to construct the Gleason score in prostate cancer, for example, are derived from separate clonal expansions of cells.

How Do Chemopreventive Drugs Slow or Stop Clonal Evolution?

This question requires a few background comments before it can be answered. Combustion products of cigarettes and fossil fuels contain both mutagenic and proliferation-inducing mitogenic molecules, as well as irritant molecules that induce epithelial proliferation associated with reactive inflammation. It is a striking fact that the number of years before lung cancer appears after starting to smoke is inversely related to the number of cigarettes smoked per day⁴⁾. Interpreted at the cellular level, this observation indicates that during the progression of intraepithelial neoplasia induced in the respiratory mucosa by concurrent exposure to mutagenic and mitogenic molecules in cigarette smoke, the more cigarettes smoked per day, the faster is the rate of clonal evolution to the point of invasive neoplasia, i.e., cancer. This example in smokers illustrates the general principle, amply confirmed by animal experiments⁵⁾, that the RATE of clonal evolution in human carcinogenesis is continuously driven by the DOSE LEVEL of concurrent exposure to mutagens and mitogens found in the environment and endogenously⁶⁾.

Thus, chemopreventive drugs work by suppressing the two concurrent driving forces of neoplastic clonal evolution: mutagenesis and mitogenesis⁷⁾. Accordingly they are categorized as being either antimutagens, antimitogens, or both.

Environmental oxidants, such as the organic oxides and peroxides found in cigarette smoke of fossil fuel exhausts, and prooxidants, which are molecules that cause increased production in cells of superoxide, hydrogen peroxide, and hydroxyl

free radicals, are both mutagenic and mitogenic. The antioxidant chemopreventive agents that block them form a category of chemopreventives that are both antimutagenic and antimitogenic.

Finally, the nonsteroidal antiinflammatory agents (NSAIDs) form a category of chemopreventives that work by blocking the arrival at an inflammatory site of monocytes, macrophages, and neutrophils producing extracellular superoxide and peroxide. NSAIDs are essentially antioxidant, and therefore are both antimutagenic and antimitogenic in their action. One example of an antimutagen is oltipraz, a dithiolethione, which induces glutathione-s-transferase in tissues, thereby accelerating the reaction of glutathione with activated carcinogens that results in their inactivation and excretion. An example of an antimitogen is tamoxafen, which blocks the stimulation of proliferation by estrogen in estrogen-sensitive tissues. Examples of antioxidants are vitamins C, E, and the many plant phenolics found in green vegetables such as the flavonoids, all of which react with oxidant molecules to form stable, non-reactive compounds that "soak up" or scavenge free radical electrons⁶⁾. Finally, aspirin is an example of a category of non-steroidal antiinflammatory agents which, as a rule, are cancer chemopreventive in animal models, and, in the case of aspirin, likely in humans as well⁸⁾.

The sharp contrast between drugs used for chemoprevention and drugs used for cancer chemotherapy should be noted. Chemoprevention uses drugs that are antimutagenic, whereas chemotherapy uses the highly promutagenic alkylating agents such as nitrogen mustards and nitrosoureas. Chemoprevention uses antioxidants, whereas chemotherapy uses highly prooxidant antibiotics, such as doxorubicin, bleomycin, and mitomycin C, which are redox cyclers that ge-

nerate mutagenic and cytotoxic free radical oxygen.

WHAT ARE SURROGATE ENDPOINT BIOMARKERS (SEBs), AND WHY ARE THEY NEEDED IN CLINICAL TRIALS OF CHEMOPREVENTIVE AGENTS?

Surrogate endpoint biomarkers (SEBs) are molecular, cellular, or tissue changes associated with the neoplastic process which occur prior to the stage of invasiveness, i.e., during the stage of intraepithelial neoplasia (see Fig. 1), and which exhibit a measurable response to chemopreventive agents that accurately predicts their effect on cancer incidence.

At present, a serious barrier to development of the field of chemoprevention is the unacceptable cost (millions of dollars), long duration (5~10 years), and large scale of effort (thousands of subjects) required by clinical trials which use the endpoint of cancer incidence reduction. It is urgent that the endpoint of cancer incidence reduction be replaced by surrogate endpoint biomarkers, which occur much earlier during the preinvasive phase of neoplastic development. With such surrogate endpoints the trials can be carried out in a year or less, and, depending on the precision of the assay for surrogate endpoints, may require fewer subjects.

WHAT IS THE ROLE OF COMPUTER-ASSISTED QUANTITATIVE IMAGE ANALYSIS (CQIA) IN IMPROVING THE PRECISION OF SEB ASSAYS?

The advent of computer-assisted cytometric techniques offers the promise of quietly revolutionizing the practice of diagnostic histopathology.

The sensitive and precise measurements of the morpho- and photometric parameters of dysplasia made by the computerized image cytometer, when compared to the present-day alternative of subjectively estimating such categories as "moderate to severe", "pleomorphism", or "hyperchromasia", offer increased diagnostic precision that is greater by many orders of magnitude.

Basically, two modalities are used in image analysis.

One, cytomorphometry, measures geometric relationships, such as nuclear dimensions, chromatin texture, and nucleolar size, shape, and position. The other, cytophotometry, measures cell and nuclear optical density at different wavelengths after staining with different dyes. Since there exist innumerable tissue surrogate biomarker assays (see below) which depend on the amount of chromogen development fixed to a cell or nucleus by chemical antibody, or cDNA probes that are marked by a second chromogen generating molecule (e.g., a fluorescent dye or horseradish peroxidase), the range of applications of image cytophotometry is very wide.

WHAT POSSIBLE CHOICES ARE THERE FOR SEBs THAT CAN BE USED IN CHEMOPREVENTION TRIALS?

In choosing a SEB for use in chemoprevention clinical trials, a variety of structural alterations characteristic of neoplasia at every level of organization is available for consideration. At the genomic level, many potentially useful SEBs are possible, based on the different types of genomic instability which occur during neoplasia (reviewed in 4). Examples are DNA single and double strand breakage processes, point mutations, mismatch repair mutations, microsatellite instability, gene

amplification and allelic loss, and the karyotypic aberrations of aneuploidy and aneusomy. Activated oncogenes and inactivated tumor suppressor genes have frequently been considered as potential SEBs in spite of their relatively poor sensitivity; they are present in about only about 50% or less of common carcinomas (p53 is an exception, occurring in up to 70% of cases of colorectal carcinoma and 90% of pancreatic carcinoma). At the cytoplasmic level, aberrantly synthesized differentiation molecules offer promise as SEBs, e.g., aberrant glycosylation of glycoproteins, including the Ta, T, and sialo-T antigens⁴⁾.

The well-known growth factors PDGF, EGF, TGF, FGF, IGF, and their receptors, are possessed by epithelial cells as well as inflammatory cells⁹⁾. Attempts to use growth factors/receptors as SEBs faces the serious problem of variable contamination of tissue preparations by inflammatory cells possessing the same growth factors/receptors, and adequate control of this variability may be difficult. At the cellular level, the proliferative index and nuclear /nucleolar morphometric parameters measured by computer-aided quantitative image analysis appear to be useful SEBs; they are discussed in more detail below.

WHAT ARE THE BEST CHOICES FOR SURROGATE ENDPOINT MARKERS?

The Chemoprevention Branch, National Cancer Institute, is now sponsoring 18 clinical trials of chemopreventive agents in breast ductal carcinoma in situ, lung dysplasia, colon adenomatous polyps, cervical intraepithelial neoplasia, bladder superficial neoplasia, and skin actinic keratoses. In each of these trials, the basic SEB used are those based on the morphologic and functional criteria used by the pathologist to make the diagnosis of

intraepithelial neoplasia, measured by computer-aided quantitative image analysis. These diagnostic criteria are: increased nuclear size, abnormal nuclear shape, increased nuclear stain uptake, pleomorphism (increased variability of nuclear size, shape, and stain uptake), increased mitoses, abnormal mitoses, and abnormal or absent maturation. In glandular epithelia such as breast and prostate, an additional criterion is the presence of an increased number of nucleoli showing enlargement, abnormal shape, and pleomorphism of size and shape. SEBs based on these criteria, and measured by computer-aided quantitative image analysis, are: proliferative index, ploidy status, nuclear morphometric parameters of increased nuclear size, altered nuclear shape, altered nuclear texture, abnormal nuclear variation in size, shape, and texture, and nucleolar morphometric parameters of nucleolar number, size, shape, position, and pleomorphism.

Proliferative index: The proliferative index is measured using antibodies against PCNA, Ki-67, Mib-1, and BrdU associated antigens, by tritiated-thymidine uptake/autoradiography, and by mitotic counts. The proliferative index has proven to be a reliable prognostic factor in breast cancer, accurately predicting the recurrence-free survival and overall survival either by itself¹⁰⁾, or as part of a commonly used grading system¹¹⁾.

Ploidy status: Aneuploidy has been shown to occur during intraepithelial neoplasia in bladder¹²⁾, prostate^{13,14)}, breast¹⁵⁻¹⁸⁾, and cervix, skin, oral leukoplakia, larynx, lung, esophagus, stomach, and colorectum (reviewed in 19). In one study of breast DCIS, the cribriform pattern exhibited 38% aneuploidy, whereas the comedo pattern exhibited 82% aneuploidy¹⁷⁾. In another study, atypical hyperplasia of the breast also exhibited aneuploidy in 4 of 13 cases¹⁸⁾. With regard to invasive

neoplasia, ploidy status has proven to be a reliable prognostic factor in both breast^{19,20)}, and prostate²¹⁾ cancer.

Nuclear morphometry (nuclear size, shape, texture, and pleomorphism). Pleomorphism is measured by the pleomorphism index, which is the sum of the CV's associated with measurements of nuclear size, shape, and texture. It is remarkable that in a number of studies, alteration of nuclear shape alone has proven to be a better predictor of mortality in stage A2 prostatic cancer than has the Gleason, Mostofi, or Johns Hopkins grading systems, or ploidy status^{22~24)}. Multivariate analysis of up to 16 nuclear shape descriptors, including nuclear roundness factor, variance of roundness factor, and nuclear ellipticity, have accurately predicted recurrence of cancer after surgery in 11 of 26 patients with renal cell carcinoma²⁵⁾, 7/14 patients with transitional cell carcinoma of the bladder²⁶⁾, and 17/27 patients with wilms' tumor (kidney)²⁷⁾.

Nucleolar morphometry (number, size, shape, position, and pleomorphism): Nucleoli are ribosome factories expressed by genes located on the acrocentric chromosomes 13, 14, 15, 21, and 22. In the activation of proliferation or secretion, the size and sometimes the number of nucleoli increases. In a study of eight nuclear and nine nucleolar morphometric features in breast cancer, simple nucleolar frequency, or the total number of nucleoli per 100 nuclei, was the best single predictor of recurrence-free survival²⁸⁾. Changes in nucleolar morphometry have been reported to be a correlate of the extent of neoplastic progression in prostatic intraepithelial neoplasia^{29,30)}. Using the silver stain for nucleolar organizing regions, the AgNOR stain, in lesions of the colorectum, the mean number of nucleolar organizer regions (NORs) clearly distinguish between tubular

adenomas, villous adenomas with moderate nuclear atypia, villous adenomas with severe nuclear atypia, and colorectal adenocarcinoma³¹⁾.

REFERENCES

- 1) Nowell PC. The clonal evolution of tumor cell populations. Acquired genetic liability permits stepwise selection of variant. sublines and underlies tumor progression. *Science* 1976; 194: 23-28.
- 2) Nowell PC. Mechanisms of tumor progression. *Cancer Res.* 1986; 46: 2203-2207.
- 3) Vogelstein B, Fearon ER, Hamilton SR, Kern SE, Preisinger AC, Leppert M, Nakamura Y, White R, Smits AMM, Bos JL. Genetic alterations during colorectal-tumor development. *N Engl J Med* 1988; 319: 525-532.
- 4) Boone CW, Kelloff GJ. Development of surrogate endpoint biomarkers for clinical trials of cancer chemopreventive agents: Relationships to fundamental properties of preinvasive (intraepithelial) neoplasia. *J Cell Biochem* 1994; 19: 10-22.
- 5) Truhaut R (ed.) In: Truhaut, R. (ed.) Potential Carcinogenic Hazards from Drugs. Evaluation of Risks. New York: Springer-Verlag, Vol. 7, 1967
- 6) Boone CW, Kelloff GJ, Freedman LS. Intraepithelial and postinvasive neoplasia as a stochastic continuum of clonal evolution, and its relationship to mechanisms of chemopreventive drug action. *J Cell Biochem* 1993; 17G: 14-25.
- 7) Boone CW, Kelloff GJ. Intraepithelial neoplasia, surrogate endpoint biomarkers, and cancer chemoprevention. *J Cell Biochem* 1993; 17F: 37-48.
- 8) Thun MJ, Namboodiri MM, Heath CW, Jr. Aspirin use and reduced risk of fatal colon cancer. *N Engl J Med* 1991; 325: 1593-1596.
- 9) Sporn MB, Roberts AB: "Peptide growth factors and their receptors" Vol 2. New York: Springer-Verlag, 1991
- 10) van Diest PJ, Baak JP, Matze-Cok P, Wisse-Breklemans EC, van Galen CM, Kurver PH, Bellot SM, Fijnheer J, van Gorp LH, Kwee WS, et al. Reproducibility of mitosis counting in 2,469 breast cancer specimens: Results from the multicenter morphometric mammary carcinoma project. *Hum Pathol* 1992; 23: 603-607.
- 11) Bloom HJG. Prognosis in carcinoma of the breast.

- Br J Cancer* 1950; 4: 259-288.
- 12) Tribukait B. DNA flow cytometry in carcinoma of the prostate for diagnosis, prognosis and study of tumor biology. *Acta Oncol* 1991; 30: 187-192.
 - 13) Montironi R, Scarpelli M, Sisti S, Braccischi A, Gusella P, Pisani E, Alberti R, Mariuzzi GM. Quantitative analysis of prostatic intraepithelial neoplasia on tissue sections. *Anal Quant Cytol Histol* 1990; 12: 366-372.
 - 14) Amin MB, Schultz DS, Zarbo RJ, Kubus J, Shaheen C. Computerized static DNA ploidy analysis of prostatic intraepithelial neoplasia. *Arch Pathol Lab Med* 1993; 117: 794-798.
 - 15) Visscher DW, Micale MA, Crissman JD. Pathological and biological relevance of cytophotometric DNA content to breast carcinoma genetic progression. *J Cell Biochem* 1993; 17G: 114-122.
 - 16) Erhardt K, Auer GU. Mammary carcinoma. Comparison of nuclear DNA content from *in situ* and infiltrative components. *Anal Quant Cytol Histol* 1987; 9: 263-267.
 - 17) Crissman JD, Visscher DW, Kubus J. Image cytophotometric DNA analysis of atypical hyperplasias and intraductal carcinomas of the breast. *Arch Pathol Lab Med* 1990; 114: 1249-1253.
 - 18) Carpenter R, Gibbs N, Matthews J, Cooke T. Importance of cellular DNA content in pre-malignant breast disease and pre-invasive carcinoma of the female breast. *Br J Surg* 1987; 74: 905-906.
 - 19) Boone CW, Kelloff GJ, Steele VE. Natural history of intraepithelial neoplasia in humans with implications for cancer chemoprevention strategy. *Cancer Res* 1992; 52: 1651-1659.
 - 20) van Dist PJ, Baak JPA. Quantitative Cyto- and Histoprognosis in Breast Cancer, pp. 55-62. New York: Elsevier, 1992
 - 21) Lieber MM. DNA ploidy: Early malignant lesions. *J Cell Biochem* 1992; 16H: 44-46.
 - 22) Epstein JI, Berry SJ, Eggleston JC. Nuclear roundness factor. A predictor of progression in untreated stage A2 prostate cancer. *Cancer* 1984; 54: 1666-1671.
 - 23) Mohler JL, Parin AW, Epstein JI, Becker RL, Mikel UV, Sesterhenn IA, Mostofi FK, Gleason DF, Sharief Y, Coffey DS. Prediction of prognosis in untreated stage A2 prostatic carcinoma. *Cancer* 1992; 69: 511-519.
 - 24) Partin AW, Walsh AC, Pitcock RV, Mohler JL, Epstein JI, Coffey DS. A comparison of nuclear morphometry and Gleason grade as a predictor of prognosis in stage A2 prostate cancer: A critical analysis. *J Urol* 1989; 142: 1254-1258.
 - 25) Pound CR, Partin AW, Epstein JI, Simons JW, Marshall FF. Nuclear morphometry accurately predicts recurrence in clinically localized renal cell carcinoma. *Urology* 1993; 42: 243-248.
 - 26) Borland RN, Partin AW, Epstein JI, Brendler CB. The use of nuclear morphometry in predicting recurrence of transitional cell carcinoma. *J Urol* 1993; 149: 272-275.
 - 27) Gearhart JP, Partin AW, Leventhal B, Beckwith JB, Epstein JI. The use of nuclear morphometry to predict response to therapy in Wilms' tumor. *Cancer* 1992; 69: 804-808.
 - 28) van Diest PJ, Mouriquand J, Schipper NW, Baak JPA. Prognostic value of nucleolar morphometric variables in cytological breast cancer specimens. *J Clin Pathol* 1990; 43: 157-159.
 - 29) Montironi R, Braccischi A, Matera G, Scarpelli M, Pisani E. Quantitation of the prostatic intra-epithelial neoplasia. Analysis of the nucleolar size, number and location. *Pathol Res Pract* 1991; 187: 307-314.
 - 30) Helpap B. Observations on the number, size and localization of nucleoli in hyperplastic and neoplastic prostatic disease. *Histopathology* 1988; 13: 203-211.
 - 31) Yang P, Huang GS, Zhu XS. Role of nucleolar organizer regions in differentiating malignant from benign tumours of the colon. *J Clin Pathol* 1990; 43: 235-238.

대한암예방학회지 : 제 2 권 제 2 호 1997

대한암예방학회지 : 제 2 권 제 2 호 1997

대한암예방학회지 : 제 2 권 제 2 호 1997

- Biomarkers of Premalignant Neoplasia and their use as Surrogate Endpoints
in Clinical Trials of Chemopreventive Agents-
- Biomarkers of Premalignant Neoplasia and their use as Surrogate Endpoints
in Clinical Trials of Chemopreventive Agents-
- Biomarkers of Premalignant Neoplasia and their use as Surrogate Endpoints
in Clinical Trials of Chemopreventive Agents-

Chares W. Boone

Chares W. Boone

Chares W. Boone